

- Porritt, G. T. (1876) Capture of Lepidoptera in East Sussex. *Entomologist's Mon. Mag.* 13:37-38.
- Robbins, R. W. (1892) Notes on collecting. *Entomologist's Rec. J. Var.* 3: 232.
- Russell, A. (1912) Seasonal notes for 1912. *Entomologist's Rec. J. Var.* 25: 248-9.
- Salvage, S. (1975) Unpublished diaries.
- Sich, A. (1912) *Euchloe cardamines* ovipositing on *Capsella bursa-pasturis*. *Entomologist's Rec. J. Var.* 24: 221.
- Southerby, R. M. (1883) Lepidoptera of Eastbourne. *Entomologist* 16:22-23.
- Thornewill, Rev. C. F. (1883) Lepidoptera of Abbot's Wood. *Entomologist* 16:35.
- Tugwell, W. H. (1892) The Lepidoptera of East Sussex in Early June. *Entomologist* 25: 156.
- Whinstone, J. P. (1883) Abbot's Wood in July. *Entomologist* 16: 183-4.
- Worsley Wood, H. (121) *Brenthis argynnis selene*, second brood at Abbot's Wood *Entomologist* 54: 17.
- 

EILEMA CANIOLA HUBN. (THE HOARY FOOTMAN) IN KENT — On the night of 23rd/24th August 1984 a specimen of *E. caniola* was captured in an actinic trap at Greatstone, Kent. According to Chalmers-Hunt (*The Butterflies and Moths of Kent* 3: 235) this represents the fifth confirmed record of this species for Kent this century. Although a number of *Autographa gamma* L. were taken on the same night, there were no other notable immigrants seen. My thanks are due to Bernard Skinner for confirming the identity of this moth. A. P. CLANCY, Delhi Cottage, Dungeness, Kent, TN29 9NE.

COPRIS LUNARIS L. (COL.: SCARABAEIDAE) ON GUERNSEY — At about 12.05 a.m. on the night of June 14th/15th 1986 a male specimen of this rare dung beetle flew to my m.v. trap at Two Wells Barn, in the Vale district of Guernsey. According to notes very kindly provided by Mr. L. Jessop of the British Museum (Natural History) this species has not been on mainland Britain since 1955, but is said to be widespread in France. I cannot find any previous reference to this species on Guernsey. My trap was sited in agricultural land, and there were tethered cattle in close proximity. *C. lunaris* excavates deep brood tunnels beneath cow or horse dung. The light, sandy soils of Vale especially on the nearby L'Ancrese Common could, I am sure, be searched profitably for the surface "spoil heaps" that indicate the underground activities of this interesting beetle. — M. D. BRYAN, Keeper of Natural History, Birmingham Museum.